

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.


Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ment APB containing the given angle. Draw the diameter PD perpendicular to AB. Also draw AQ perpendicular to AB. Draw BP meeting AQ in Q. With B as a center, and a radius equal to p, describe an arc cutting AQ produced in R. With R as a center, and a radius equal to p+AQ, describe an arc cutting AB produced in S. On SR measure off SN=to SA. Then RN=the altitude. Take AL=RN, and draw



*HLC* parallel to *AB*, cutting the circle in *C*. Draw *AC*, *BC*. Then *ACB* is the required triangle. For let x, y, z be the sides *BC*, *AC*, and the altitude. Then  $xy\sin C=az$ , x+y+z=p,  $a^2=x^2+y^2-2xy\cos C$ .

$$\therefore a^2 + 2xy(1 + \cos A) = (p-z)^2. \quad \therefore z^2 - 2(p + a\cot \frac{1}{2}C) = a^2 - p^2.$$

$$\therefore z = p + a \cot \frac{1}{2}C - \sqrt{(p + a \cot \frac{1}{2}C)^2 - (p^2 - a^2)}$$
.

$$\angle AQB = \frac{1}{2}C$$
,  $AQ = a\cot\frac{1}{2}C$ ,  $RS = p + a\cot\frac{1}{2}C$ ,  $AR = \sqrt{(p^2 - a^2)}$ ,  $AS = \sqrt{(p + a\cot\frac{1}{2}C)^2 - (p^2 - a^2)}$ ].

 $\therefore RN=z$ .  $\therefore$  The triangle ACB contains all the required parts.

Also solved by J. Scheffer.

## 353. Proposed by L. H. McDONALD, M. A., Ph. D., Sometimes Tutor at Cambridge, Jersey City, N. J.

In a given circle place two chords which shall be in a given ratio and also a given distance apart.

## Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Let the ratio m:n to distance apart=d; the radius of the given circle =r; u, v the distances of the chords from the center.

Then  $2\sqrt{(r^2-u^2)}$ ,  $2\sqrt{(r^2-v^2)}$  are the lengths of the chords.

$$\therefore m_1 / (r^2 - v^2) = n_1 / (r^2 - u^2)$$
, or  $(m^2 - n^2)r^2 = m^2v^2 - n^2u^2$ , and  $u + v = d$ .

$$\therefore u = \frac{m^2d - \sqrt{[r^2(m^2 - n^2)^2 + m^2n^2d^2]}}{m^2 - n^2},$$

$$v = \frac{1/[r^2(m^2-n^2)^2+m^2n^2d^2]-n^2d}{m^2-n^2}$$
.

Hence, if AB=d, take AC=u, CB=v, and with C as center and radius r describe a circle, through A and B perpendicular to AB draw lines intersecting this circle. The chords of the circle formed by these lines are the chords required.

Also solved by S. A. Corey.

## 354. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

Find the condition that triangles which are circumscribed to one of two confocal parabolas may be inscribed in the other.